

Coseparable corings

J. Gómez-Torrecillas

Departamento de Álgebra
Facultad de Ciencias
Universidad de Granada
E18071 Granada, Spain
e-mail: torrecil@ugr.es

A. Louly

Département de Mathématiques
Faculté de Sciences
Université Abdelmalek Esaadi
B.P. 2121 Tétouan, Morocco

Introduction

M. Sweedler [12] introduced the notion of a coring as a generalization of the concept of coalgebra in order to study the intermediate division rings for an extension of division rings. It turns out that this formalism embodies several kinds of relative module categories. Thus, graded modules, Doi-Hopf modules and, more generally, entwined modules are instances of comodules over suitable corings (see [1] and its references). This explains the increasing interest in the study of corings, as one could expect that the development of the theory of corings will contribute significantly to the understanding of the aforementioned relative modules.

Coseparable corings were investigated from a (co)homological point of view by F. Guzman [8], dualizing earlier results by M. Kleiner in the framework of rings [9], and extending Y. Doi's theory on coseparable coalgebras [5]. In this paper, we will relate coseparability and semisimplicity in the framework of coring theory, proving the following theorem.

Theorem. *Let A be a separable algebra over a field k . The following conditions are equivalent for an A -coring $\mathfrak{C}$.*

- (i) $\mathfrak{C}$ is coseparable;
- (ii) $\mathfrak{C} \otimes_k K$ is a semisimple $A \otimes_k K$ -coring for every field extension $k \subseteq K$;
- (iii) $\mathfrak{C} \otimes_k \mathfrak{D}$ is a semisimple $A \otimes_k B$ -coring for every semisimple B -coring $\mathfrak{D}$ (B is any k -algebra);
- (iv) $\mathfrak{C} \otimes_k \mathfrak{C}^\circ$ is a semisimple $A \otimes_k A^\circ$ -coring.

When particularized to coalgebras over fields, this theorem gives [5, Proposition 12] and [3, Theorem 3.4]. The notions involved in the statements of our theorem, like semisimple coring, opposite coring, or tensor product of corings, are defined below in this note.

1 Bicomodules

We first recall from [12] the notion of a coring. The concepts of comodule and bicomodule over a coring are generalizations of the corresponding notions for coalgebras, and were considered in [8].

Throughout this paper, $A, A', \dots, B, \dots$ denote associative and unitary algebras over a commutative ring k . The tensor product over A is denoted by $\otimes_A$. We shall sometimes replace $\otimes_k$ by $\otimes$.

1.1. Corings. An A -coring is a three-tuple $(\mathfrak{C}, \Delta_{\mathfrak{C}}, \epsilon_{\mathfrak{C}})$ consisting of an A -bimodule $\mathfrak{C}$ and two A -bimodule maps

$$\Delta_{\mathfrak{C}} : \mathfrak{C} \longrightarrow \mathfrak{C} \otimes_A \mathfrak{C} \quad \epsilon_{\mathfrak{C}} : \mathfrak{C} \longrightarrow A \quad (1)$$

such that the diagrams

$$\begin{array}{ccc} \mathfrak{C} & \xrightarrow{\Delta_{\mathfrak{C}}} & \mathfrak{C} \otimes_A \mathfrak{C} \\ \Delta_{\mathfrak{C}} \downarrow & & \downarrow \mathfrak{C} \otimes_A \Delta_{\mathfrak{C}} \\ \mathfrak{C} \otimes_A \mathfrak{C} & \xrightarrow{\Delta_{\mathfrak{C}} \otimes_A \mathfrak{C}} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \mathfrak{C} \end{array} \quad (2)$$

and

$$\begin{array}{ccc} \mathfrak{C} & \xrightarrow{\Delta_{\mathfrak{C}}} & \mathfrak{C} \otimes_A \mathfrak{C} \\ \searrow \cong & & \downarrow \mathfrak{C} \otimes_A \epsilon_{\mathfrak{C}} \\ & & \mathfrak{C} \otimes_A A \end{array} \quad \begin{array}{ccc} \mathfrak{C} & \xrightarrow{\Delta_{\mathfrak{C}}} & \mathfrak{C} \otimes_A \mathfrak{C} \\ \searrow \cong & & \downarrow \epsilon_{\mathfrak{C}} \otimes_A \mathfrak{C} \\ & & A \otimes_A \mathfrak{C} \end{array} \quad (3)$$

commute.

From now on, $\mathfrak{C}, \mathfrak{C}', \dots, \mathfrak{D}, \dots$ will denote corings over $A, A', \dots, B, \dots$, respectively.

1.2. Comodules. A *right $\mathfrak{C}$ -comodule* is a pair (M, ρ_M) consisting of a right A -module M and an A -linear map $\rho_M : M \rightarrow M \otimes_A \mathfrak{C}$ such that the diagrams

$$\begin{array}{ccc} M & \xrightarrow{\rho_M} & M \otimes_A \mathfrak{C} \\ \rho_M \downarrow & & \downarrow M \otimes_A \Delta_{\mathfrak{C}} \\ M \otimes_A \mathfrak{C} & \xrightarrow{\rho_M \otimes_A \mathfrak{C}} & M \otimes_A \mathfrak{C} \otimes_A \mathfrak{C} \end{array} \quad \begin{array}{ccc} M & \xrightarrow{\rho_M} & M \otimes_A \mathfrak{C} \\ \searrow \cong & & \downarrow M \otimes_A \epsilon_{\mathfrak{C}} \\ & & M \otimes_A A \end{array} \quad (4)$$

commute. Left $\mathfrak{C}$ -comodules are similarly defined; we use the notation λ_M for their structure maps. A *morphism* of right $\mathfrak{C}$ -comodules (M, ρ_M) and (N, ρ_N) is an A -linear map $f : M \rightarrow N$ such that the following diagram is commutative.

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \rho_M \downarrow & & \downarrow \rho_N \\ M \otimes_A \mathfrak{C} & \xrightarrow{f \otimes_A \mathfrak{C}} & N \otimes_A \mathfrak{C} \end{array} \quad (5)$$

The k -module of all right $\mathfrak{C}$ -comodule morphisms from M to N is denoted by $\text{Hom}_{\mathfrak{C}}(M, N)$. The k -linear category of all right $\mathfrak{C}$ -comodules will be denoted by $\mathcal{M}^{\mathfrak{C}}$. When $\mathfrak{C} = A$, with the trivial coring structure, the category $\mathcal{M}^A$ is just the category of all right A -modules, which is ‘traditionally’ denoted by $\mathcal{M}_A$.

Coproducts and cokernels in $\mathcal{M}^{\mathfrak{C}}$ do exist and can be already computed in $\mathcal{M}_A$. Therefore, $\mathcal{M}^{\mathfrak{C}}$ has arbitrary inductive limits. If ${}_A\mathfrak{C}$ is a flat module, then $\mathcal{M}^{\mathfrak{C}}$ is easily proved to be an abelian category.

1.3. Bicomodules. Let $\rho_M : M \rightarrow M \otimes_A \mathfrak{C}$ be a comodule structure over an $A' - A$ -bimodule M , and assume that ρ_M is A' -linear. For any right A' -module X , the right A -linear map $X \otimes_{A'} \rho_M : X \otimes_{A'} M \rightarrow X \otimes_{A'} M \otimes_A \mathfrak{C}$ makes $X \otimes_{A'} M$ a right $\mathfrak{C}$ -comodule. This leads to an additive functor $- \otimes_{A'} M : \mathcal{M}_{A'} \rightarrow \mathcal{M}^{\mathfrak{C}}$. When $A' = A$ and $M = \mathfrak{C}$, the functor $- \otimes_A \mathfrak{C}$ is right adjoint to the forgetful functor $U_A : \mathcal{M}^{\mathfrak{C}} \rightarrow \mathcal{M}_A$ (see [8, Proposition 3.1], [1, Lemma 3.1]). Now assume that the $A' - A$ -bimodule M is also a left $\mathfrak{C}'$ -comodule with structure map $\lambda_M : M \rightarrow \mathfrak{C}' \otimes_A M$. It is clear that $\rho_M : M \rightarrow M \otimes_A \mathfrak{C}$ is a morphism of left $\mathfrak{C}'$ -comodules if and only if $\lambda_M : M \rightarrow \mathfrak{C}' \otimes_{A'} M$ is a morphism of right $\mathfrak{C}$ -comodules. In this case, we say that M is a $\mathfrak{C}' - \mathfrak{C}$ -bicomodule. The $\mathfrak{C}' - \mathfrak{C}$ -bicomodules are the objects of a k -linear category ${}^{\mathfrak{C}'}\mathcal{M}^{\mathfrak{C}}$ whose morphisms are those $A' - A$ -bimodule maps which are morphisms of $\mathfrak{C}'$ -comodules and of $\mathfrak{C}$ -comodules. Some particular cases are now of interest. For instance, when $\mathfrak{C}' = A'$, the objects of the category ${}^{A'}\mathcal{M}^{\mathfrak{C}}$ are the $A' - A$ -bimodules with a right $\mathfrak{C}$ -comodule structure $\rho_M : M \rightarrow M \otimes_A \mathfrak{C}$ which is A' -linear.

We will use the following lemma.

1.4. Lemma. *If M and N are A -bimodules, and P and L are B -bimodules, then*

$$(M \otimes_A N) \otimes_k (L \otimes_B P) \cong (M \otimes_k L) \otimes_{A \otimes_k B} (N \otimes_k P),$$

and this is a natural isomorphism of $A \otimes_k B$ -bimodules.

Proof. Straightforward. □

With the aid of this lemma, we can give the tensor product of corings as follows.

1.5. Proposition. *Let $(\mathfrak{C}, \Delta_{\mathfrak{C}}, \epsilon_{\mathfrak{C}})$ and $(\mathfrak{D}, \Delta_{\mathfrak{D}}, \epsilon_{\mathfrak{D}})$ be corings over the algebras A and B , respectively. Then $\mathfrak{C} \otimes_k \mathfrak{D}$ is an $A \otimes_k B$ -coring, with the comultiplication*

$$\mathfrak{C} \otimes_k \mathfrak{D} \xrightarrow{\Delta_{\mathfrak{C}} \otimes_k \Delta_{\mathfrak{D}}} (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes_k (\mathfrak{D} \otimes_B \mathfrak{D}) \cong (\mathfrak{C} \otimes_k \mathfrak{D}) \otimes_{A \otimes_k B} (\mathfrak{C} \otimes_k \mathfrak{D}),$$

and the counit

$$\mathfrak{C} \otimes_k \mathfrak{D} \xrightarrow{\epsilon_{\mathfrak{C}} \otimes_k \epsilon_{\mathfrak{D}}} A \otimes_k B$$

Proof. Clearly, the proposed comultiplication and counit are homomorphisms of $A \otimes_k B$ -bimodules. To check the coassociative property, consider the diagram

$$\begin{array}{ccccc}
& & (\mathfrak{C} \otimes \mathfrak{C}) \otimes (\mathfrak{D} \otimes \mathfrak{D}) & & \\
& \nearrow \Delta \otimes \Delta & \downarrow & \searrow \cong & \\
\mathfrak{C} \otimes \mathfrak{D} & & (\mathfrak{C} \otimes_A \Delta) \otimes (\mathfrak{D} \otimes_B \Delta) & & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) \\
& \downarrow \Delta \otimes \Delta & & & \downarrow (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\Delta \otimes \Delta) \\
(\mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D}) & & & & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D}) \\
& \searrow (\Delta \otimes \mathfrak{C}) \otimes (\Delta \otimes \mathfrak{D}) & \downarrow & \nearrow \cong & \downarrow \cong \\
& & (\mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D} \otimes_B \mathfrak{D}) & & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) \\
& \downarrow \cong & \downarrow \cong & & \downarrow \cong \\
& & (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) & & \\
& \nearrow (\Delta \otimes \Delta) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) & & \nearrow \cong & \\
& & (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) & &
\end{array}$$

whose commutativity follows from that of the inner diagrams. The counitary property follows analogously from the commutativity of the diagram

$$\begin{array}{ccc}
\mathfrak{C} \otimes \mathfrak{D} & \xrightarrow{\Delta \otimes \Delta} & (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes (\mathfrak{D} \otimes_B \mathfrak{D}) \xrightarrow{\cong} (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) \\
& \searrow \cong & \downarrow (\mathfrak{C} \otimes \epsilon) \otimes (\mathfrak{D} \otimes \epsilon) \quad \downarrow (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\epsilon \otimes \epsilon) \\
& & (\mathfrak{C} \otimes_A A) \otimes (\mathfrak{D} \otimes_B A) \xrightarrow{\cong} (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (A \otimes B)
\end{array}$$

□

1.6. Example. One important source of examples of corings are the entwining structures, introduced by T. Brzeziński and S. Majid in [2]. Let A, C be a k -algebra and a k -coalgebra, respectively. Assume that $A \otimes C$ is endowed with an A -bimodule structure, where the left A -module structure is the canonical one. The right A -module structure induces a k -linear map $\psi : C \otimes A \rightarrow A \otimes C$ given by $\psi(c \otimes a) = (1 \otimes c)a$. Consider the maps

$$A \otimes C \xrightarrow{A \otimes \Delta_C} A \otimes C \otimes C \cong (A \otimes C) \otimes_A A \otimes C ,$$

and

$$A \otimes C \xrightarrow{A\epsilon_C} A \otimes k \cong A.$$

By [1, Proposition 2.2] they give a structure of A -coring on $A \otimes C$ if and only if (A, C, ψ) is an *entwining structure* over k . The tensor product of corings given in Proposition 1.5 provides a canonical definition of tensor product of entwining structures as follows. Let (B, D, φ) be a second entwining structure over k , and consider the tensor product $A \otimes B$ -coring $A \otimes C \otimes B \otimes D$. The isomorphism of k -modules

$$A \otimes C \otimes B \otimes D \cong A \otimes B \otimes C \otimes D \quad (6)$$

allows to induce an $A \otimes B$ -bimodule structure on $A \otimes B \otimes C \otimes D$ from the bimodule structure of the $A \otimes B$ -coring $A \otimes C \otimes B \otimes D$. The resulting left $A \otimes B$ -module structure is then the canonical one, while the right $A \otimes B$ structure is, after some computations, the induced by the map

$$C \otimes D \otimes A \otimes B \cong C \otimes A \otimes D \otimes B \xrightarrow{\psi \otimes \varphi} A \otimes C \otimes B \otimes D \cong A \otimes B \otimes C \otimes D,$$

which will be denoted by $\psi \otimes \varphi$. It can be checked in a straightforward way that the comultiplication and the counit of the tensor product $A \otimes B$ -coring $A \otimes C \otimes B \otimes D$ define, by using the isomorphism (6), a comultiplication and a counit on $A \otimes B \otimes C \otimes D$ in such a way that $(A \otimes B, C \otimes D, \psi \otimes \varphi)$ is an entwining structure, which we call *tensor product entwining structure*. Of course, (6) becomes an isomorphism of $A \otimes B$ -corings.

1.7. Opposite coring. The notation A° stands for the opposite algebra of A ; every $A - B$ -bimodule M gives in an obvious way a $B^\circ - A^\circ$ bimodule, denoted by M° . So every A -coring $\mathfrak{C}$ defines an A° -coring $\mathfrak{C}^\circ$ as follows: we have an isomorphism of k -modules $\tau : \mathfrak{C} \otimes_A \mathfrak{C} \rightarrow \mathfrak{C}^\circ \otimes_{A^\circ} \mathfrak{C}^\circ$ defined by $\tau(x \otimes_A y) = y \otimes_{A^\circ} x$. Then the k -linear map

$$\mathfrak{C} \xrightarrow{\Delta} \mathfrak{C} \otimes_A \mathfrak{C} \xrightarrow{\tau} \mathfrak{C}^\circ \otimes_{A^\circ} \mathfrak{C}^\circ$$

can be viewed as an $A^\circ - A^\circ$ -bimodule map $\Delta^\circ : \mathfrak{C}^\circ \rightarrow \mathfrak{C}^\circ \otimes_{A^\circ} \mathfrak{C}^\circ$, which is clearly coassociative. Consider the counit $\epsilon : \mathfrak{C} \rightarrow A$ as an $A^\circ - A^\circ$ -bimodule map $\epsilon^\circ : \mathfrak{C}^\circ \rightarrow A^\circ$ to obtain an A° -coring $(\mathfrak{C}^\circ, \Delta^\circ, \epsilon^\circ)$ called the *opposite coring* of $\mathfrak{C}$.

We are now in position to show that bicomodules are in fact comodules.

1.8. Proposition. *Let A, B be algebras, and $\mathfrak{C}, \mathfrak{D}$ be corings over A and B , respectively. Then there is an isomorphism of categories ${}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}} \cong {}^{\mathfrak{C}^\circ}_k \mathfrak{D}^\circ \mathcal{M}$.*

Proof. If (M, λ_M, ρ_M) is a $\mathfrak{C} - \mathfrak{D}^\circ$ -bicomodule, then the homomorphism of $A \otimes_k B$ -bimodules

$$\lambda : M \xrightarrow{\lambda_M} \mathfrak{C} \otimes_A M \xrightarrow{\mathfrak{C} \otimes_A \rho_M} \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \xrightarrow{\cong} (\mathfrak{C} \otimes_k \mathfrak{D}) \otimes_{A \otimes_k B} M$$

[illegible]
$$\begin{aligned}
(\Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ)(\mathfrak{C} \otimes_A \rho_M) \lambda_M &= (\Delta \otimes_A \rho_M \otimes_{B^\circ} \mathfrak{D}^\circ)(\mathfrak{C} \otimes_A \rho_M) \lambda_M \\
&= (\Delta \otimes_A \rho_M \otimes_{B^\circ} \mathfrak{D}^\circ)(\lambda_M \otimes_{B^\circ} \mathfrak{D}) \rho_M \\
&= (\mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \rho_M \otimes_{B^\circ} \mathfrak{D}^\circ)(\Delta \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ)(\lambda_M \otimes_{B^\circ} \mathfrak{D}) \rho_M \\
&= (\mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \rho_M \otimes_{B^\circ} \mathfrak{D}^\circ)(\mathfrak{C} \otimes_A \lambda_M \otimes_{B^\circ} \mathfrak{D}^\circ)(\mathfrak{C} \otimes_A \rho_M) \lambda_M
\end{aligned}$$
$$\begin{array}{ccccccc}
M & \xrightarrow{\lambda_M} & \mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A^{\rho_M}} & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \xrightarrow{\cong} & (\mathfrak{C} \otimes_k \mathfrak{D}) \otimes_{A \otimes B} M \\
& \searrow \cong & \downarrow \epsilon \otimes_A M & & \downarrow \epsilon \otimes_A M \otimes_{B^\circ} \epsilon^\circ & & \downarrow (\epsilon \otimes \epsilon) \otimes_{A \otimes B} M \\
& & A \otimes_A M & \xrightarrow{\cong} & A \otimes_A M \otimes_{B^\circ} & \xrightarrow{\cong} & (A \otimes B) \otimes_{A \otimes B} M
\end{array}$$

is also commutative, the counital property for the coaction λ holds, and we have that (M, λ) is a left $\mathfrak{C}$ -comodule. Moreover, it is easy to see that every homomorphism of $\mathfrak{C} - \mathfrak{D}^\circ$ -bicomodules becomes a morphism of left $\mathfrak{C} \otimes_k \mathfrak{D}$ -comodules.

Conversely, let (M, λ) any left $\mathfrak{C} \otimes_k \mathfrak{D}$ -comodule. Then the commutativity of the diagrams

$$\begin{array}{c}
 \begin{array}{ccccc}
 M & & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
 \lambda \searrow & & \cong \nearrow & & \searrow \mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ \\
 & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & & \Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ & \\
 \downarrow \lambda & & \downarrow \Delta \otimes_A M & & \downarrow \mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ \\
 (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & \xrightarrow{(\mathfrak{C} \otimes \mathfrak{D}) \otimes \lambda} & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & \xrightarrow{\cong} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \mathfrak{D}^\circ \\
 \downarrow \cong & & \downarrow \cong & & \downarrow \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ \otimes_{B^\circ} \epsilon^\circ \\
 \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \xrightarrow{\mathfrak{C} \otimes \lambda \otimes \mathfrak{D}^\circ} & \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M) \otimes_{B^\circ} \mathfrak{D}^\circ & \xrightarrow{\cong} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \mathfrak{D}^\circ \\
 \downarrow \mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ & & \downarrow \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M) \otimes_{B^\circ} \epsilon^\circ & & \downarrow \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ \otimes_{B^\circ} \epsilon^\circ \\
 \mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes \lambda} & \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M) & \xrightarrow{\cong} & (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \\
 & & & & \downarrow (\mathfrak{C} \otimes_A \mathfrak{C}) \otimes_A M \otimes_{B^\circ} \epsilon^\circ \\
 & & & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M
 \end{array}
 \end{array}$$

and

$$\begin{array}{ccccc}
 M & \xrightarrow{\lambda} & (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & \xrightarrow{\cong} & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \xrightarrow{\mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ} \mathfrak{C} \otimes_A M \\
 & \searrow \cong & \downarrow (\epsilon \otimes \epsilon) \otimes_{A \otimes B} M & & \downarrow \epsilon \otimes_A M \otimes_{B^\circ} \epsilon^\circ \\
 & & (A \otimes B) \otimes_{A \otimes B} M & \xrightarrow{\cong} & A \otimes_A M \otimes_{B^\circ} B^\circ \xrightarrow{\cong} A \otimes_A M
 \end{array}$$

show that (M, λ_M) is a left $\mathfrak{C}$ -comodule with structure map

$$\lambda_M : M \xrightarrow{\lambda} (\mathfrak{C} \otimes_k \mathfrak{D}) \otimes_{A \otimes_k B} M \xrightarrow{\cong} \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \xrightarrow{\mathfrak{C} \otimes_A M \otimes_{B^\circ} \epsilon^\circ} \mathfrak{C} \otimes_A M.$$

Analogously, the map

$$\rho_M : M \xrightarrow{\lambda} (\mathfrak{C} \otimes_k \mathfrak{D}) \otimes_{A \otimes_k B} M \xrightarrow{\cong} \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \xrightarrow{\epsilon \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ} M \otimes_{B^\circ} \mathfrak{D}^\circ$$

endows M with a structure of a right $\mathfrak{D}^\circ$ -comodule. Next, we shall check that (M, λ_M, ρ_M) is a $\mathfrak{C} - \mathfrak{D}^\circ$ -bicomodule. Consider for this the commutative diagrams

$$\begin{array}{c}
\begin{array}{ccccc}
M & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & & \\
\lambda \searrow & & \cong \nearrow & & \Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ \searrow \\
& (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \mathfrak{D}^\circ \\
\lambda \downarrow & & \Delta \otimes_{A \otimes B} M & & \downarrow \\
(\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & & & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \epsilon^\circ \\
& \searrow (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} \lambda & & \Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ & \\
& (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} (\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M & & & \downarrow \\
& \downarrow & & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \\
& \cong & & & \downarrow \\
& \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& \searrow \mathfrak{C} \otimes_A \lambda \otimes_{B^\circ} \mathfrak{D}^\circ & & \cong \nearrow & \\
& \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M) \otimes_{B^\circ} \mathfrak{D}^\circ & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ \otimes_{B^\circ} \epsilon^\circ & \\
& \downarrow & & \downarrow & \\
& \mathfrak{C} \otimes_A M & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& \searrow \mathfrak{C} \otimes_A \lambda & & \cong \nearrow & \\
& \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathfrak{D}) \otimes_{A \otimes B} M) & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& & & \searrow \mathfrak{C} \otimes_A \epsilon \otimes M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& & & \downarrow & \\
& & & \mathfrak{C} \otimes_A \epsilon \otimes M \otimes_{B^\circ} \mathfrak{D}^\circ & \\
& & & \downarrow & \\
& & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathfrak{D}^\circ &
\end{array}
\end{array}$$

and

$$\begin{array}{c}
\begin{array}{ccccc}
M & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ & & \\
\lambda \searrow & & \cong \nearrow & & \Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ \searrow \\
& (\mathfrak{C} \otimes \mathcal{D}) \otimes_{A \otimes B} M & & & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ \otimes_{B^\circ} \mathcal{D}^\circ \\
\lambda \downarrow & & & & \downarrow \\
(\mathfrak{C} \otimes \mathcal{D}) \otimes_{A \otimes B} M & \xrightarrow{(\mathfrak{C} \otimes \mathcal{D}) \otimes \lambda} & (\mathfrak{C} \otimes \mathcal{D}) \otimes_{A \otimes B} M & \xrightarrow{\Delta \otimes_A M \otimes_{B^\circ} \Delta^\circ} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ \otimes_{B^\circ} \mathcal{D}^\circ \\
\cong \downarrow & & \downarrow & & \downarrow \\
\mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ & \xrightarrow{\mathfrak{C} \otimes \lambda \otimes \mathcal{D}^\circ} & \mathfrak{C} \otimes_A ((\mathfrak{C} \otimes \mathcal{D}) \otimes_{A \otimes B} M) \otimes_{B^\circ} \mathcal{D}^\circ & \xrightarrow{\cong} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ \otimes_{B^\circ} \mathcal{D}^\circ \\
\epsilon \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ \downarrow & & \downarrow & & \downarrow \\
M \otimes_{B^\circ} \mathcal{D}^\circ & \xrightarrow{\lambda \otimes \mathcal{D}^\circ} & ((\mathfrak{C} \otimes \mathcal{D}) \otimes_{A \otimes B} M) \otimes_{B^\circ} \mathcal{D}^\circ & \xrightarrow{\cong} & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ \otimes_{B^\circ} \mathcal{D}^\circ \\
& & \downarrow & & \downarrow \\
& & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ & & \mathfrak{C} \otimes_A M \otimes_{B^\circ} \mathcal{D}^\circ
\end{array}
\end{array}$$

which say that $(\mathfrak{C} \otimes_A \rho_M) \lambda_M = (\lambda_M \otimes_{B^\circ} \mathcal{D}^\circ) \rho_M$, that is, M is a bicomodule. Moreover, it is easy to see that every homomorphism of left $\mathfrak{C} \otimes_k \mathcal{D}$ -comodules is a homomorphism of $\mathfrak{C} - \mathcal{D}^\circ$ -bicomodules. We have constructed two functors which are easily seen to be mutually inverse. This gives the desired isomorphism of categories. $\square$

2 Coseparable corings and semisimple corings

Recall from [4] that the algebra A over the commutative ring k is said to be *separable* if the multiplication map $\mu : A \otimes_k A \rightarrow A$ splits as an A -bimodule epimorphism, i.e., there is a homomorphism of A -bimodules $\theta : A \rightarrow A \otimes_k A$ such that $\mu\theta = \text{id}_A$. Dually, one obtains the notion of a coseparable coalgebra and, more generally, of a coseparable coring. Following [8], we say that an A -coring $\mathfrak{C}$ is *coseparable* if there is a $\mathfrak{C}$ -bicomodule map $\pi : \mathfrak{C} \otimes_A \mathfrak{C} \rightarrow \mathfrak{C}$ such that $\pi\Delta = \text{id}_{\mathfrak{C}}$ (here, A is not assumed to be separable).

Both notions can be expressed functorially by means of the separable functors introduced by C. Năstăsescu, M. Van den Bergh and F. Van Oystaeyen in [10]. Thus, A is a

separable k -algebra if and only if the forgetful functor $\mathcal{M}_A \rightarrow \mathcal{M}_k$ is separable [10, Proposition 1.3], and $\mathfrak{C}$ is a coseparable coring if and only if the forgetful functor $\mathcal{M}^{\mathfrak{C}} \rightarrow \mathcal{M}_A$ is separable [1, Corollary 3.6]. We take advantage of this functorial approach in our development (see [7] for a general treatment of separable functors for corings). Separable functors are nicely characterized when they are in adjunction, see [11, Theorem 1.2]. The first theorem in this section needs a previous lemma.

2.1. Lemma. *Let $\mathfrak{C}$ and $\mathfrak{D}$ be corings over k -algebras A and B , respectively. If $F : {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}} \rightarrow {}^A\mathcal{M}^{\mathfrak{D}}$ and $F' : {}^A\mathcal{M}^{\mathfrak{D}} \rightarrow \mathcal{M}^{\mathfrak{D}}$ are the forgetful functors, then*

- (1) *The functor $G = \mathfrak{C} \otimes_A - : {}^A\mathcal{M}^{\mathfrak{D}} \rightarrow {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}}$ is right adjoint to the functor F with unit $\lambda : 1 \rightarrow GF$ given at every $M \in {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}}$ by the left $\mathfrak{C}$ -comodule structure map $\lambda_M : M \rightarrow \mathfrak{C} \otimes_A M$.*
- (2) *The functor $G' = A \otimes_k - : \mathcal{M}^{\mathfrak{D}} \rightarrow {}^A\mathcal{M}^{\mathfrak{D}}$ is left adjoint to F' with the unit $c : G'F' \rightarrow 1$ defined at every $M \in {}^A\mathcal{M}^{\mathfrak{D}}$ by the left A -module structure map $c_M : A \otimes_k M \rightarrow M$.*

Proof. (1) This is [8, Proposition 3.1].

(2) Straightforward. □

2.2. Theorem. *Let $\mathfrak{C}$ be a coseparable A -coring, and assume that A is a separable algebra over the commutative ring k . If $\mathfrak{D}$ is a coring over a k -algebra B , then the forgetful functor $U : {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}} \rightarrow \mathcal{M}^{\mathfrak{D}}$ is separable.*

Proof. We first decompose the forgetful functor U accordingly the following diagram

$$\begin{array}{ccc} {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}} & \xrightarrow{U} & \mathcal{M}^{\mathfrak{D}} \\ & \searrow F \quad \nearrow F' & \\ & {}^A\mathcal{M}^{\mathfrak{D}} & \end{array} .$$

We shall prove that F and F' are separable functors, which implies, by [10, Lemma 1.1], that U is separable. By [11, Theorem 1.2], F is separable if and only if there is a natural transformation $\mu : GF \rightarrow 1$ such that $\mu \circ \lambda = 1_{{}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}}}$. By [1, 3.6], there is a morphism of A -bimodules $\gamma : \mathfrak{C} \otimes_A \mathfrak{C} \rightarrow A$ such that $\gamma \Delta = \epsilon$ and $(\mathfrak{C} \otimes_A \gamma)(\Delta \otimes_A \mathfrak{C}) = (\gamma \otimes_A \mathfrak{C})(\mathfrak{C} \otimes_A \Delta)$. Then, for every $\mathfrak{C} - \mathfrak{D}$ -bicomodule (M, λ_M, ρ_M) , the commutativity of the following diagrams

$$\begin{array}{ccccc} \mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A \lambda_M} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M & \xrightarrow{\gamma \otimes_A M} & M \\ \mathfrak{C} \otimes_A \rho_M \downarrow & & \downarrow \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \rho_M & & \downarrow \rho_M \\ \mathfrak{C} \otimes_A M \otimes_B \mathfrak{D} & \xrightarrow{\mathfrak{C} \otimes_A \lambda_M \otimes_B \mathfrak{D}} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \otimes_B \mathfrak{D} & \xrightarrow{\gamma \otimes_A M \otimes_B \mathfrak{D}} & M \otimes_B \mathfrak{D} \end{array}$$

$$\begin{array}{ccccc}
\mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A \lambda_M} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M & \xrightarrow{\gamma \otimes_A M} & M \\
\Delta \otimes_A M \downarrow & & & & \downarrow \lambda_M \\
\mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \lambda_M} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A \gamma \otimes_A M} & \mathfrak{C} \otimes_A M
\end{array}$$

gives that $\mu_M : \mathfrak{C} \otimes_A M \xrightarrow{\mathfrak{C} \otimes_A \lambda_M} \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M \xrightarrow{\gamma \otimes_A M} M$ is a homomorphism of $\mathfrak{C}$ – $\mathfrak{D}$ -bicomodules. On the other hand, if $f : M \rightarrow N$ is a homomorphism of $\mathfrak{C}$ – $\mathfrak{D}$ -bicomodules, then the diagram

$$\begin{array}{ccccc}
\mathfrak{C} \otimes_A M & \xrightarrow{\mathfrak{C} \otimes_A \lambda_M} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A M & \xrightarrow{\gamma \otimes_A M} & M \\
\mathfrak{C} \otimes_A f \downarrow & & & & \downarrow f \\
\mathfrak{C} \otimes_A N & \xrightarrow{\mathfrak{C} \otimes_A \lambda_N} & \mathfrak{C} \otimes_A \mathfrak{C} \otimes_A N & \xrightarrow{\gamma \otimes_A N} & N
\end{array}$$

is commutative. Therefore we have defined a natural transformation $\mu : GF \rightarrow 1_{\mathfrak{C}\mathcal{M}^{\mathfrak{D}}}$. Moreover, for every $M \in \mathfrak{C}\mathcal{M}^{\mathfrak{D}}$, one has $\mu_M \circ \lambda_M = (\gamma \otimes_A M)(\Delta \otimes_A M)\lambda_M = (\epsilon \otimes_A M)\lambda_M = M$. Thus, λ splits off, and F is a separable functor.

Now, to prove that F' is a separable functor it suffices, by [11, Theorem 1.2], to show that there is a natural transformation $\nu : 1 \rightarrow G'F'$ such that $c \circ \nu = 1$. Since A is a separable k -algebra, we have an A -bimodule map $\theta : A \rightarrow A \otimes_k A$ such that $\mu\theta = A$, where $\mu : A \otimes_k A \rightarrow A$ is the multiplication map. Then for every A – $\mathfrak{D}$ -bicomodule M we have the commutative diagram

$$\begin{array}{ccccccc}
M & \xrightarrow{\cong} & A \otimes_A M & \xrightarrow{\theta \otimes_A M} & A \otimes A \otimes_A M & \xrightarrow{\cong} & A \otimes M \\
\rho_M \downarrow & & \downarrow A \otimes_A \rho_M & & \downarrow A \otimes A \otimes_A \rho_M & & \downarrow A \otimes \rho_M \\
M \otimes_B \mathfrak{D} & \xrightarrow{\cong} & A \otimes_A M \otimes_B \mathfrak{D} & \xrightarrow{\theta \otimes_A M \otimes_B \mathfrak{D}} & A \otimes A \otimes_A M \otimes_B \mathfrak{D} & \xrightarrow{\cong} & A \otimes M \otimes_B \mathfrak{D},
\end{array}$$

which says that $\nu_M : M \xrightarrow{\cong} A \otimes_A M \xrightarrow{\theta \otimes_A M} A \otimes_k A \otimes_A M \xrightarrow{\cong} A \otimes_k M$ is $\mathfrak{D}$ -colinear and, thus, a homomorphism of A – $\mathfrak{D}$ -bicomodules. On the other hand, if $f : M \rightarrow N$ is homomorphism of A – $\mathfrak{D}$ -bicomodules, then we have the commutativity of the following diagram

$$\begin{array}{ccccccc}
M & \xrightarrow{\cong} & A \otimes_A M & \xrightarrow{\theta \otimes_A M} & A \otimes A \otimes_A M & \xrightarrow{\cong} & A \otimes M \\
f \downarrow & & & & & & \downarrow A \otimes f \\
N & \xrightarrow{\cong} & A \otimes_A N & \xrightarrow{\theta \otimes_A N} & A \otimes A \otimes_A N & \xrightarrow{\cong} & A \otimes N.
\end{array}$$

Thus we have defined a natural transformation $\nu : 1_{A\mathcal{M}^{\mathfrak{D}}} \rightarrow G'F'$. Moreover, for every $M \in {}^A\mathcal{M}^{\mathfrak{D}}$, one has $c_M\nu_M = M$. This proves that F' is separable, and the theorem holds. $\square$

An A -coring $\mathfrak{C}$ is called *semisimple* if $\mathfrak{C}$ satisfies one of the equivalent conditions in the following Theorem 2.3. Here, a comodule is said to be simple if it has just two subcomodules. A semisimple comodule is a direct sum of simple comodules.

2.3. Theorem ([6]). *Let $\mathfrak{C}$ be an A -coring. The following statements are equivalent:*

- (i) $\mathfrak{C}$ is semisimple as a left $\mathfrak{C}$ -comodule and $\mathfrak{C}_A$ is flat;
- (ii) every left $\mathfrak{C}$ -comodule is semisimple and $\mathfrak{C}_A$ is flat;
- (iii) $\mathfrak{C}$ is semisimple as a right $\mathfrak{C}$ -comodule and ${}_A\mathfrak{C}$ is flat;
- (iv) every right $\mathfrak{C}$ -comodule is semisimple and ${}_A\mathfrak{C}$ is flat;
- (v) every (left or right) $\mathfrak{C}$ -comodule is semisimple, and ${}_A\mathfrak{C}$ and $\mathfrak{C}_A$ are projectives.

Coseparable corings over separable algebras can be characterized in terms of semisimplicity. The first step to obtain the characterization is the following.

2.4. Theorem. *Let $\mathfrak{C}$ be a coseparable A -coring such that ${}_A\mathfrak{C}$ is flat. If A is a separable k -algebra, then the $A \otimes_k B$ -coring $\mathfrak{C} \otimes_k \mathfrak{D}$ is semisimple for every semisimple B -coring $\mathfrak{D}$.*

Proof. Since ${}_A\mathfrak{C}$ and ${}_B\mathfrak{D}$ are flat, we get that $\mathfrak{C} \otimes_k \mathfrak{D}$ is flat as a left $A \otimes_k B$ -module. By Proposition 1.8, we have an isomorphism of categories ${}^{\mathfrak{C} \otimes_k \mathfrak{D}}\mathcal{M} \cong {}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}^\circ}$. By Theorem 2.2, the forgetful functor ${}^{\mathfrak{C}}\mathcal{M}^{\mathfrak{D}^\circ} \rightarrow \mathcal{M}^{\mathfrak{D}^\circ} \cong {}^{\mathfrak{D}}\mathcal{M}$ is separable. Therefore we obtain a separable functor ${}^{\mathfrak{C} \otimes_k \mathfrak{D}}\mathcal{M} \rightarrow {}^{\mathfrak{D}}\mathcal{M}$. Now, $\mathfrak{D}$ is assumed to be semisimple, which implies, by [10, Proposition 1.2], that $\mathfrak{C} \otimes_k \mathfrak{D}$ is semisimple. $\square$

2.5. Remark. The $A \otimes_k B$ -coring $\mathfrak{C} \otimes_k \mathfrak{D}$ fails to be semisimple if A is not assumed to be separable. For example, the trivial structure of A -coring given on A by the canonical isomorphism $A \cong A \otimes_A A$ is obviously coseparable. Assume k to be a field. If A is not semisimple as a ring, then $A \cong A \otimes_k k$ is not semisimple as a $A \otimes_k k \cong A$ -coring (but k is obviously a semisimple k -coring).

Our next goal is to obtain a characterization of coseparable corings over separable algebras over a field that generalizes the given for separable coalgebras in [3, Theorem 3.4]. To achieve this, we need to prove some results which are also of independent interest.

Let us start with a generalization of the so called fundamental theorem of coalgebras. The notion which replaces the finite dimensionality over fields is given by the following definition.

2.6. Definition. An $R - S$ -bimodule M is said to be of *finite type* if it is a finitely generated left $R \otimes_k S^\circ$ -module, that is, if $M = \sum_i Rm_iS$, for some $m_i \in M$, where i runs over a finite set.

Following [12], let us endow the left dual ${}^*\mathfrak{C} = \text{Hom}_A(\mathfrak{C}, {}_A A)$ with a structure of ring by the defining its multiplication by $gf = g(\mathfrak{C} \otimes_A f)\Delta$. The map $A^\circ \rightarrow {}^*\mathfrak{C}$ induced by the counit $\epsilon : \mathfrak{C} \rightarrow A$ is then a ring homomorphism. Analogously, the right dual $\mathfrak{C}^* = \text{Hom}_A(\mathfrak{C}, A_A)$ is a ring with multiplication $gf = f(g \otimes_A \mathfrak{C})\Delta$, and we have a ring homomorphism $A^\circ \rightarrow \mathfrak{C}^*$. Then the obvious structure of A° -bimodule of $\mathfrak{C}$ extends to a ${}^*\mathfrak{C} - \mathfrak{C}^*$ -bimodule structure. Here, the left ${}^*\mathfrak{C}$ -action is given by $f.c = (f \otimes_A \mathfrak{C})\Delta$, and the right $\mathfrak{C}^*$ -action is given by $c.f = (\mathfrak{C} \otimes_A f)\Delta$.

2.7. Proposition. *Let $\mathfrak{C}$ be an A -coring such that ${}_A \mathfrak{C}$ and $\mathfrak{C}_A$ are projective modules. Then $\mathfrak{C} = \cup_i \mathfrak{C}_i$, where the $\mathfrak{C}_i$'s are $\mathfrak{C}$ -subbicomodules of $\mathfrak{C}$ which are of finite type as A -bimodules.*

Proof. Thinking of $\mathfrak{C}$ as a left ${}^*\mathfrak{C} \otimes_k \mathfrak{C}^*$ -module, we can write it as a direct union of its finitely generated submodules. Thus, $\mathfrak{C} = \cup_i \mathfrak{C}_i$, where the $\mathfrak{C}_i$ are ${}^*\mathfrak{C} - \mathfrak{C}^*$ -subbimodules of $\mathfrak{C}$ of finite type. Let us prove that each $\mathfrak{C}_i$ is of finite type as an A -bimodule. By [6, Corollary 2.11], the $\mathfrak{C}$ -subbicomodules of $\mathfrak{C}$ are precisely its ${}^*\mathfrak{C} - \mathfrak{C}^*$ -subbimodules. Thus, accordingly with [6, Proposition 2.10], this proof is finished as soon as we prove that every rational ${}^*\mathfrak{C} - \mathfrak{C}^*$ -bimodule of finite type M is of finite type as an A -bimodule. For, if $M = \sum_s {}^*\mathfrak{C} m_s \mathfrak{C}^*$, for finitely many $m_s \in M$, then we proceed as follows: choose a system of left rational parameters (c_{sj}, m_{sj}) for each m_s (see [6] for this notion), and then choose a system of right rational parameters (m_{sjk}, c_{sjk}) for every m_{sj} . If $m \in M$, then $m = \sum_s b_s m_s b'_s$, for some $b_s \in {}^*\mathfrak{C}, b'_s \in \mathfrak{C}^*$. Compute

$$m = \sum_{s,j} (m_{sj} b_s(c_{sj})) b'_s = \sum_{s,j} (m_{sj} b'_s) b_s(c_{sj}) = \sum_{s,j,k} b'_s(c_{sjk}) m_{sjk} b_s(c_{sj}),$$

whence $M = \sum_{s,j,k} A m_{sjk} A$, and it is an A -bimodule of finite type. This finishes our proof. $\square$

A coring $\mathfrak{C}$ is said to be *simple* if it has no nontrivial subbicomodules. Examples of simple A -corings are the Sweedler's canonical corings $A \otimes_B A$, where B is a simple artinian subring of A (see [6, Proposition 4.2]).

2.8. Corollary. *If $\mathfrak{C}$ is a simple coring such that ${}_A \mathfrak{C}$ and $\mathfrak{C}_A$ are projective modules, then $\mathfrak{C}$ is an A -bimodule of finite type.*

Every homomorphism of A -corings $\varphi : \mathfrak{C} \rightarrow \mathfrak{D}$ induces a functor $(-)_\varphi : {}^\mathfrak{C} \mathcal{M} \rightarrow {}^\mathfrak{D} \mathcal{M}$, called corestriction functor. The corresponding homomorphism of rings $\varphi^* : \mathfrak{D}^* \rightarrow \mathfrak{C}^*$ (see [12, Proposition 3.2]) gives the usual restriction of scalars functor $(-)_\varphi^* : \mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{\mathfrak{D}^*}$.

2.9. Proposition. *Let $\varphi : \mathfrak{C} \rightarrow \mathfrak{D}$ be a homomorphism of A -corings. If $\mathfrak{C}_A$ and $\mathfrak{D}_A$ are finitely generated projective modules, then the following conditions are equivalent:*

- (i) *The functor $(-)_\varphi : {}^\mathfrak{C} \mathcal{M} \rightarrow {}^\mathfrak{D} \mathcal{M}$ is separable;*
- (ii) *the functor $(-)_\varphi^* : \mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{\mathfrak{D}^*}$ is separable.*

Proof. By [1, Lemma 4.3], we have isomorphisms of categories ${}^{\mathfrak{C}}\mathcal{M} \cong \mathcal{M}_{\mathfrak{C}^*}$ and ${}^{\mathfrak{D}}\mathcal{M} \cong \mathcal{M}_{\mathfrak{D}^*}$. We get the result from the commutativity of the following diagram of functors

$$\begin{array}{ccc} {}^{\mathfrak{C}}\mathcal{M} & \xrightarrow{(-)_{\varphi}} & {}^{\mathfrak{D}}\mathcal{M} \\ \downarrow \cong & & \downarrow \cong \\ \mathcal{M}_{\mathfrak{C}^*} & \xrightarrow{(-)_{\varphi^*}} & \mathcal{M}_{\mathfrak{D}^*}. \end{array}$$

□

2.10. Theorem. *Let $\mathfrak{C}$ be a coring over a separable k -algebra A . If $\mathfrak{C}_A$ is projective and finitely generated then $\mathfrak{C}$ is a coseparable coring if and only if $\mathfrak{C}^*$ is a separable k -algebra.*

Proof. Consider the counit $\epsilon : \mathfrak{C} \rightarrow A$ as a homomorphism of corings. Then the functor $(-)_{\epsilon} : {}^{\mathfrak{C}}\mathcal{M} \rightarrow {}_A\mathcal{M}$ is just the forgetful functor, and, in this way, $(-)_{\epsilon}$ is separable if and only if $\mathfrak{C}$ is a coseparable coring. On the other hand, the functor $(-)_{\epsilon^*} : \mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{A^{\circ}}$ is nothing but the restriction of scalars functor associated to the ring homomorphism $\epsilon^* : A^{\circ} \rightarrow \mathfrak{C}^*$. Thus, by Proposition 2.9, $\mathfrak{C}$ is coseparable if and only if $\mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{A^{\circ}}$ is separable. On the other hand, by [10, Proposition 1.3], we have that the restriction of scalars functor $\mathcal{M}_{A^{\circ}} \rightarrow \mathcal{M}_k$ is separable, since A° is a separable k -algebra. Now, the composite $\mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{A^{\circ}} \rightarrow \mathcal{M}_k$ is the evident restriction of scalars functor. Hence, by [10, Lemma 1.1], this functor is separable if and only if $\mathcal{M}_{\mathfrak{C}^*} \rightarrow \mathcal{M}_{A^{\circ}}$ is. In other words, $\mathfrak{C}$ is a coseparable coring if and only if $\mathfrak{C}^*$ is a separable k -algebra. □

Let R be a ring and M a right R -module. Recall that an R -submodule N of M is said to be *pure* if the canonical map $N \otimes_R L \rightarrow M \otimes_R L$ is injective for every left R -module L . Pure submodules of left R -modules are similarly defined. We say that a homomorphism of right R -modules $f : X \rightarrow Y$ is pure if $\text{Im} f$ is a pure R -submodule of Y .

2.11. Proposition. *Let $\varphi : \mathfrak{C} \rightarrow \mathfrak{D}$ be a homomorphism of A -corings. If φ is a pure monomorphism of right A -modules, then the functor $(-)_{\varphi} : {}^{\mathfrak{C}}\mathcal{M} \rightarrow {}^{\mathfrak{D}}\mathcal{M}$ is separable.*

Proof. Let $M, N \in {}^{\mathfrak{C}}\mathcal{M}$. If $f : M \rightarrow N$ is a homomorphism of $\mathfrak{D}$ -comodules, then it is a homomorphism of $\mathfrak{C}$ -comodules. In fact, one has

$$(\varphi \otimes_A N) \lambda_N f = (\mathfrak{D} \otimes_A f)(\varphi \otimes_A M) \lambda_M = (\varphi \otimes_A N)(\mathfrak{C} \otimes_A f) \lambda_M.$$

Since φ is a pure monomorphism of right A -modules, we get that $\lambda_N f = (\mathfrak{C} \otimes_A f) \lambda_M$. Therefore we can define the map $\varphi_{M,N} : \text{Hom}_{\mathfrak{D}}(M, N) \rightarrow \text{Hom}_{\mathfrak{C}}(M, N)$ by $\varphi_{M,N}(f) = f$. Thus, by definition (see [10, p. 398]), $(-)_{\varphi}$ is a separable functor. □

A subbicomodule $\mathfrak{D}$ of a coring $\mathfrak{C}$ is said to be a *subcoring* if $\mathfrak{D}$ is pure in $\mathfrak{C}$ both as a left and as a right A -submodule. In such a case, the restriction of the comultiplication of $\mathfrak{C}$ to $\mathfrak{D}$ gives a well defined structure of A -coring on $\mathfrak{D}$ such that the inclusion $\mathfrak{D} \subseteq \mathfrak{C}$ is a homomorphism of A -corings. Every subbicomodule of a semisimple coring is a subcoring, since it is a direct summand as a left and as a right A -submodule. The following is a consequence of Proposition 2.11 and [10, Proposition 1.2].

2.12. Corollary. *Every subcoring of a semisimple coring is semisimple.*

The following result tell us that the separability of a morphism of corings defined on a semisimple coring is a ‘local’ property.

2.13. Proposition. *Let $\mathfrak{C}$ be a semisimple A -coring and let $\varphi : \mathfrak{C} \rightarrow \mathfrak{D}$ be a homomorphism of A -corings such that the image of φ is pure in $\mathfrak{D}$ both as a left and right A -module. Then $(-)_\varphi : {}^{\mathfrak{C}}\mathcal{M} \rightarrow {}^{\mathfrak{D}}\mathcal{M}$ is a separable functor if and only if the functor $(-)_{\varphi|_{\mathfrak{C}'}} : {}^{\mathfrak{C}'}\mathcal{M} \rightarrow {}^{\mathfrak{D}'}\mathcal{M}$ is separable for every pair of subcorings $\mathfrak{C}' \leq \mathfrak{C}$ and $\mathfrak{D}' \leq \mathfrak{D}$ of finite type as A -bimodules such that $\varphi(\mathfrak{C}') \subseteq \mathfrak{D}'$.*

Proof. Assume $(-)_\varphi$ to be a separable functor, and let $\mathfrak{C}'$ and $\mathfrak{D}'$ as declared. If $i : \mathfrak{C}' \rightarrow \mathfrak{C}$ and $j : \mathfrak{D}' \rightarrow \mathfrak{D}$ denote the inclusion maps, then $(-)_j \circ (-)_{\varphi|_{\mathfrak{C}'}} = (-)_\varphi \circ (-)_i$, which is separable by Proposition 2.11. Apply [10, Lemma 1.1] to obtain that $(-)_{\varphi|_{\mathfrak{C}'}}$ is separable. Conversely, since $\mathfrak{C}$ is semisimple, we get from [6, Theorem 3.5] a decomposition $\mathfrak{C} = \bigoplus_i \mathfrak{C}_i$, with $\mathfrak{C}_i$ a simple semisimple A -coring for every i . By 2.8, all the $\mathfrak{C}_i$ ’s are A -bimodules of finite type. Write $\mathfrak{D}_i = \varphi(\mathfrak{C}_i)$, which is pure in $\mathfrak{D}$ as a left and as a right A -submodule for every i . Therefore, these $\mathfrak{D}_i$ ’s are subcorings of $\mathfrak{D}$ which are of finite type as A -bimodules. Let us denote by $\varphi_i : \mathfrak{C}_i \rightarrow \mathfrak{D}_i$ the coring homomorphisms induced by φ . By hypothesis, the functors $(-)_{\varphi_i}$ are separable which implies, by [7, Theorem 4.7], that there exists bicomodule homomorphisms $\omega_i : \mathfrak{C}_i \square_{\mathfrak{D}_i} \mathfrak{C}_i \rightarrow \mathfrak{C}_i$ such that $\omega_i \overline{\Delta_{\mathfrak{C}_i}} = \mathfrak{C}_i$ for every i . Here, $\overline{\Delta_{\mathfrak{C}_i}}$ is given by the comultiplication on $\mathfrak{C}_i$. Finally, $\mathfrak{C}_i \square_{\mathfrak{D}_i} \mathfrak{C}_i = \mathfrak{C}_i \square_{\mathfrak{D}} \mathfrak{C}_i$ and $\mathfrak{C} \square_{\mathfrak{D}} \mathfrak{C} = \bigoplus_i (\mathfrak{C}_i \square_{\mathfrak{D}} \mathfrak{C}_i) \oplus \bigoplus_{i \neq j} \mathfrak{C}_i \square_{\mathfrak{D}} \mathfrak{C}_j$. Define $\omega = \bigoplus_i \omega_i$ over $\bigoplus_i (\mathfrak{C}_i \square_{\mathfrak{D}} \mathfrak{C}_i)$ and $\omega = 0$ over $\bigoplus_{i \neq j} \mathfrak{C}_i \square_{\mathfrak{D}} \mathfrak{C}_j$; we have $\omega \overline{\Delta} = \mathfrak{C}$. By [7, Theorem 4.7], $(-)_\varphi$ is a separable functor. $\square$

If we apply Proposition 2.13 to the homomorphism of corings given by the counity $\epsilon : \mathfrak{C} \rightarrow A$, then we obtain the following.

2.14. Corollary. *A semisimple A -coring is coseparable if and only every subcoring of finite type as A -bimodule is coseparable.*

Let $\mathfrak{C}$ be an A -coring and K a commutative ring containing k . Then $\mathfrak{C} \otimes_k K$ is a coring over $A \otimes_k K$, with left dual ring ${}^*(\mathfrak{C} \otimes_k K) = \text{Hom}_{A \otimes_k K}(\mathfrak{C} \otimes_k K, A \otimes_k K)$ endowed with the convolution product. On the other hand, we have the tensor product structure of ring on ${}^*\mathfrak{C} \otimes_k K$. These two k -algebras are related.

2.15. Proposition. *The homomorphism of k -modules*

$$\Psi : \text{Hom}_A(\mathfrak{C}, A) \otimes_k K \rightarrow \text{Hom}_{A \otimes_k K}(\mathfrak{C} \otimes_k K, A \otimes_k K)$$

defined by $\Psi(f \otimes_k \alpha)(x \otimes_k \beta) = f(x) \otimes_k \alpha \beta$ for every $f \in \text{Hom}_A(\mathfrak{C}, A) \otimes_k K$, $x \in \mathfrak{C}$, $\alpha, \beta \in K$ is a homomorphism of k -algebras. Moreover, if ${}_A \mathfrak{C}$ is a finitely generated projective module, then Ψ is an isomorphism.

Proof. Let us prove that Ψ is a homomorphism of algebras. Let $f, f' \in \text{Hom}_A(\mathfrak{C}, A)$, $x \in \mathfrak{C}$ and $\alpha, \alpha', \beta \in K$. If $\Delta(x) = x_1 \otimes_k x_2$, then

$$\begin{aligned} \Psi[(f \otimes_k \alpha)(f' \otimes_k \alpha')](x \otimes_k \beta) &= f f'(x) \otimes_k \alpha \alpha' \beta \\ &= f(x_1 f'(x_2)) \otimes_k \alpha \alpha' \beta \\ &= \Psi(f \otimes_k \alpha)[x_1 f'(x_2) \otimes_k \beta \alpha'] \\ &= \Psi(f \otimes_k \alpha)[(x_1 \otimes_k \beta) \Psi(f' \otimes_k \alpha')(x_2 \otimes_k 1_k)] \\ &= [\Psi(f \otimes_k \alpha) \Psi(f' \otimes_k \alpha')](x \otimes_k \beta). \end{aligned}$$

Now assume that ${}_A \mathfrak{C}$ is a finitely generated projective module. Let $\Phi : \text{Hom}_{A \otimes_k K}(\mathfrak{C} \otimes_k K, A \otimes_k K) \rightarrow \text{Hom}_A(\mathfrak{C}, A) \otimes_k K$ the map defined as follows: First, take a dual basis $(x_i, \varphi_i)_i$ for ${}_A \mathfrak{C}$. If $g \in \text{Hom}_{A \otimes_k K}(\mathfrak{C} \otimes_k K, A \otimes_k K)$, write $g(x_i \otimes_k 1) = \sum_j (a_{ij} \otimes_k \alpha_{ij})$, and define $\Phi(g) = \sum_{i,j} \varphi_i a_{ij} \otimes_k \alpha_{ij}$. It is not hard to see that Φ is the inverse map for Ψ . $\square$

We are now in position to prove our main result.

2.16. Theorem. *Let A be a separable algebra over a field k . The following conditions are equivalent for an A -coring $\mathfrak{C}$.*

- (i) $\mathfrak{C}$ is coseparable;
- (ii) $\mathfrak{C} \otimes_k K$ is a semisimple $A \otimes_k K$ -coring for every field extension $k \subseteq K$;
- (iii) $\mathfrak{C} \otimes_k \mathfrak{D}$ is a semisimple $A \otimes_k B$ -coring for every semisimple B -coring $\mathfrak{D}$ (B is any k -algebra);
- (iv) $\mathfrak{C} \otimes_k \mathfrak{C}^\circ$ is a semisimple $A \otimes_k A^\circ$ -coring.

Proof. First, we recall that our separable algebra A over the field k is necessarily a finite dimensional semisimple k -algebra.

(i) $\Rightarrow$ (iii) This is a particular case of Theorem 2.4.

(iii) $\Rightarrow$ (ii) Put $B = \mathfrak{D} = K$.

(ii) $\Rightarrow$ (i) The coring $\mathfrak{C}$ is semisimple, so by Corollary 2.14 and Corollary 2.12 we can assume that $\mathfrak{C}$ is an A -bimodule of finite type. This implies that $\mathfrak{C}$ is finite dimensional as a k -vector space and, in particular, ${}_A \mathfrak{C}$ is finitely generated and projective. By Proposition 2.15, $*\mathfrak{C} \otimes_k K \cong *(\mathfrak{C} \otimes_k K)$ for every field extension $k \subseteq K$. This last k -algebra is semisimple because the $A \otimes_k K$ -coring $\mathfrak{C} \otimes_k K$ is semisimple, and $\mathfrak{C} \otimes_k K$ is finitely generated and projective as a left $A \otimes_k K$ -module. By a classical result ([4, Theorem 2.5]), $*\mathfrak{C}$ becomes a separable k -algebra which implies, by Theorem 2.10, that $\mathfrak{C}$ is a coseparable coring.

(iii) $\Rightarrow$ (iv) Taking $\mathfrak{D} = B = k$ we obtain that $\mathfrak{C}$ is semisimple. By Theorem 2.3, $\mathfrak{C}^\circ$ is also semisimple. Therefore, $\mathfrak{C} \otimes_k \mathfrak{C}^\circ$ is semisimple.

(iv) $\Rightarrow$ (i) The comultiplication $\Delta : \mathfrak{C} \rightarrow \mathfrak{C} \otimes_A \mathfrak{C}$ is a homomorphism of $\mathfrak{C}$ -bicomodules. By Proposition 1.8, it is a homomorphism of left $\mathfrak{C} \otimes_k \mathfrak{C}^\circ$ -comodules. Since this coring is assumed to be semisimple, Δ splits as a left $\mathfrak{C} \otimes_k \mathfrak{C}^\circ$ -comodule map and, hence, as a $\mathfrak{C}$ -bicomodule map. Therefore, $\mathfrak{C}$ is coseparable. $\square$

References

- [1] T. Brzeziński, *The structure of corings: Induction functors, Maschke-type theorem, and Frobenius and Galois-type properties*, preprint arXiv:math.RA/0002105 v2, 2000.
- [2] T. Brzeziński and S. Majid, *Coalgebra bundles*, Commun. Math. Phys. **191** (1998), 467–492.
- [3] F. Castaño Iglesias, J. Gómez-Torrecillas, and C. Năstăsescu, *Separable functors in coalgebras. Applications.*, Tsukuba J. Math. **21** (1997), 329–344.
- [4] F. DeMeyer and E. Ingraham, *Separable Algebras over Commutative Rings*, Lecture Notes in Mathematics, vol. 181, Springer-Verlag, Berlin, 1971.
- [5] Y. Doi, *Homological coalgebra*, J. Math. Soc. Japan **33** (1981), 31–50.
- [6] L. El Kaoutit, J. Gómez-Torrecillas, and F. J. Lobillo, *Semisimple corings*, preprint arXiv:math.RA/0201070; MPS:Pure mathematics/0202009, 2001.
- [7] J. Gómez-Torrecillas, *Separable functors in corings*, preprint arXiv:math.RA/0102120; MPS:Pure mathematics/0111016, 2001.
- [8] F. Guzman, *Cointegrations, relative cohomology for comodules, and coseparable corings*, J. Algebra **126** (1989), 211–224.
- [9] M. Kleiner, *Integrations and cohomology theory*, J. Pure Appl. Algebra **1** (1985), 71–86.
- [10] C. Năstăsescu, M. Van den Bergh, and F. Van Oystaeyen, *Separable functors applied to graded rings*, J. Algebra **123** (1989), 397–413.
- [11] M. D. Rafael, *Separable functors revisited*, Comm. Algebra **18** (1990), 1445–1459.
- [12] M. Sweedler, *The predual theorem to the Jacobson-Bourbaki Theorem*, Trans. Amer. Math. Soc. **213** (1975), 391–406.